

Effect of Oat Forthe Treatment of Hypertriglyceridemia; Randomized Controlled Trial

Francisco Hernández-Pérez¹, Elizabeth Corona-Tlatelpa² and Emma Beatriz Medina-Cazares³

Abstract

Introduction: In Mexico, the cardiovascular diseases constitute the first cause of death, from which 56% are from Coronary artery disease, having a very stretch relation with Dyslipidemia.

Objective: To determine if the oat extract is effective, safe, and well tolerated for the treatment of hypertriglyceridemia.

Material and methods: Randomized clinical trial, in volunteers with serum triglyceride levels of 200 mg/dL or more. Two treatments were distributed; 40 gr. of oat, compared to 40 gr. of flaxseed in the morning fasting in both groups, with a follow-up of four weeks with baseline measurements and final lipid profile with diet control and exercise.

Analysis of the results: Student's t-test, was used for the comparison of means, the Wilcoxon signed-rank test was used for skewed populations. Fisher's exact test was used to compare percentages. It was considered a statistical significance of $p < 0.05$.

Results: Thirty volunteers were studied, 15 for the oat group and 15 for the flaxseed group; two individuals from the oat group were eliminated due to the presence of minor adverse effects, leaving only 13 volunteers.

No significant differences were found when comparing the oat group against the flaxseed group in the serum triglyceride levels ($p=0.247$ and 0.436 at the beginning and the end, respectively), no significant difference was found in the serum triglyceride levels in the oat group (initial 298.80 ± 114.60 against 307.20 ± 126.95 at the end, $p=0.798$), similar results were found in the flaxseed group (initial 351.61 ± 155.17 against 303.23 ± 117.40 at the end, $p=0.152$). It was found that in both groups there was significant reduction in weight.

Conclusions: the use of oats and flaxseeds were ineffective in lowering the serum triglyceride levels.

Keywords: Oats; Flaxseeds; Hypertriglyceridaemia; Hypercholesterolaemia

- 1 Center for Educational Research and Teacher Training IMSS, Teacher Education, Master in Clinical Research, Mexico
- 2 Resident Physician Family Medicine, UMF 31 Iztapalapa, Mexico
- 3 Nutritionist, General Hospital No. 47 Iztapalapa, Mexico

Corresponding author: Francisco Hernández-Pérez

✉ turco168@hotmail.com

Cerro del Coyote Street Manzana 11 Lote 14 Col. San Nicolás II CP 14734 DF Tlalpan, Distrito Federal, Mexico.

Tel: 56141662

Citation: Hernández-Pérez F, Tlatelpa EC, Cazares EBM. Effect of Oat Forthe Treatment of Hypertriglyceridemia; Randomized Controlled Trial. Herb Med. 2017, 3:1.

Received: March 01, 2017; **Accepted:** March 13, 2017; **Published:** March 18, 2017

Introduction

Dyslipidemia has become a growing public health problem. The overall prevalence of hypercholesterolemia in Mexico is high (23.6%), however there are studies in health institutions in which the prevalence is higher —between 35 and 48%. Where we can see part of the magnitude of the problem, which increases if other factors associated with overweight, hypertension, diabetes and smoking are added [1-3].

As a general rule, it is considered that hyperlipoproteinemia exists when plasma cholesterol is > 5.2 mmol/L (200 mg/dL) or the level of triglycerides exceeds 2.2 mmol/L (150 mg/dL). It has been found that controlling the levels of serum lipid stabilizes the atheromatous plaque inducing regression of the lesion and slowing down the development of vascular obstruction, with beneficial effects in reducing vascular events as well as related mortality; namely, persistent vascular lesions match with sustained high triglycerides levels [4-8].

It is not yet entirely clear whether hypertriglyceridemia causes heart disease or is merely a marker of it. The fact is that hypertriglyceridemia is closely correlated with the presence of two atherogenic elements which are: the presence of small, dense c-LDL particles and the reduction of the component HDL of cholesterol-HDL [9].

There are three interconnected pathways of metabolism of lipoproteins: one is the transportation of diet or exogenous fat; the second is the hepatic transport or endogenous fat; and the last one is the transport of cholesterol. These pathways are separate but the alteration of one affects the function of the other. A diet low in total fat and saturated fat in patients with coronary heart disease decreases the amount of exogenous lipoproteins and chylomicron particles, but both may be altering the liver metabolism as well as its transport, which require more than diet to modify the mortality [10-12].

Within the dietary management of the disease dietary fiber is included as a major component and within these the oatmeal stands out, which is a very common and inexpensive cereal. This cereal is high in soluble fiber and produces decreased levels of total cholesterol with a cardiovascular benefit, with an improved digestion and metabolism as well [13]. The hypocholesterolemic effect of soluble fibers appears to be related to its viscosity; thus, the fibers that form low viscosity aqueous solutions have demonstrated their ability to reduce cholesterol levels. Apparently, the compound to which is attributed the effect of decreasing serum levels of triglycerides are the beta-glucans [14-21].

The main mechanism of action of beta-glucan is due to its ability to bind bile acids, which not only increases the removal of cholesterol via feces, but its concentration in the blood is reduced, because the liver has to use it to make more bile salts that are essential for digestion. Further, by forming a physical barrier and cover the walls of the intestine it also reduces the absorption of dietary cholesterol. It also decreases hepatic synthesis of cholesterol, which contributes to lower the levels of (cholesterol) in the blood. Others suggest that the mechanism is due to increased viscosity of the bolus in the small intestine because the fiber leads to the formation of an impenetrable layer together with water in the mucosa which can act as a mechanical barrier that reduces absorption of nutrients including glucose and fatty acids [22].

In 2010 a study was conducted by Ruiz et al. where the use of oats in rats decreased cholesterol by 52.47% and triglycerides by 50.20%; however, human trails with oats have shown a decrease in cholesterol levels but so far, its effect on triglycerides is doubtful, although well tolerated without significant adverse reactions [23-30].

Materials and Methods

A randomized clinical trial was conducted, using nonprobabilistic sampling to test the efficacy, safety and tolerability of oats. Two groups were integrated: group 1 which received 40 grams of oats* and group 2 received 40 grams of flaxseed**, both groups received it milled, in fasting, dissolved in 250 ml of water, for 30 days in the mornings.

Both treatments were prepared by nutrition staff and administered to the volunteers who came to the consulting room of nutrition and weight control of the General Hospital of Zone no. 47 (HGZ 47, in Spanish) during the period from August 1 to November 30, 2014.

In all volunteers, basal serum levels of triglycerides, total cholesterol, HDL and LDL were measured at baseline and 30 days to assess efficacy of both treatments. Serum transaminases, bilirubin, urea and creatinine levels were also measured to assess the safety of both treatments.

The selection criteria were: both sexes aged between 18 and 50 years, with serum triglyceride levels equal to or greater than 200 mg/dl, which were not under professional medical or dietary treatment to lower lipid levels. All signed informed consent.

It was considered effective if serum triglyceride levels fell 16% or more of their baseline, success was also considered if cholesterol and LDL levels decreased 16% or more.

* Oats: Oat flakes, Kosland brand; manually crushed, 50% dietary fiber.

** Flaxseed: flaxseed flakes, Flax seed brand; manually crushed, 50% dietary fiber.

It was accepted as safe for both treatments if basal liver function had a maximum increase of 20% from the high levels considered normal. Confounding variables were controlled as is the diet and the intensity of physical activity.

Statistical Analysis

To calculate the sample size and for the randomization method, EPIDAT 3.1 statistical software was used.

The variable categories were compared using Fisher's exact test, given the sample size. Student t test was used to compare means between groups but to compare the effectiveness of treatments it was found that both groups had a skewed distribution so the Wilcoxon rank test was used for related tests. It was considered at $p \leq 0.05$ as statistically significant.

Results

Thirty volunteers were studied, 15 in group 1 (oats) and 15 in group 2 (flaxseed), the predominant gender was male with a total of 21 patients and 9 female patients, with an average age of 34 years: 19 (63%) had a history of smoking, 17 (56.6%) with a history of hypertriglyceridemia, and only 8 (26.6%) with hypercholesterolemia as background. The 63% of volunteers were overweight or obese in some degree.

The average of serum triglyceride levels was 326.8 mg/dL at baseline and 283.10 mg/dL at the end, while the levels of cholesterol were 209.00 mg/dL at baseline and 194.07 mg/dL at the end.

Of the initial 30 volunteers, 2 patients in group 2 (flaxseed) had an adverse reaction characterized by abdominal pain, bowel sounds and mild diarrhea persisting for more than three days so the withdrawal of both patients from the study was necessary (Figure 1).




Figure 1 Sample selection.

In the inferential analysis group 1 to group 2 were compared, no significant differences between the two groups in terms of general features were found. Age and gender were similar, as well as background; only patients in group 1 (oats) had more years of being hypertensive patients ($p=0.040$) (Tables 1 and 2).

Efficiency

When comparing the averages of initial and end lipids among the oat group of against the flaxseed group no significant differences were found; also, when treatment alone was compared, no significant difference between before and after treatment with oats nor flaxseed, considering that they were not effective (Tables 3 and 4).

Security

Both treatments were very safe as the liver and kidney constants were unchanged at the beginning or end of the study (Tables 5 and 6). It was only found that the final urea levels in the oat group was significantly higher ($p=0.041$) but in all cases without clinical significance since despite the differences, these remained within normal parameters.

Table 1 Comparison of general characteristics among the group of oats and flaxseed group.

Background	Oats	Flaxseed	p*
Age (years)	36.76 +/- 8.54	36.60 +/- 5.88	0.064
Weight (kg)	85.20 +/- 18.40	89.76 +/- 13.80	0.113
SBP (mmHg)	110.00 +/- 8.45	110.67 +/- 7.03	0.864
DBP (mmHg)	74.00 +/- 7.35	72.67 +/- 4.57	0.019
BMI	31.42 +/- 5.47	32.26 +/- 5.13	0.352
Years of diabetes	0.27 +/-1.03	0.53 +/-2.06	0.344
Years of hypertension	0.77 +/-0.25	0.0 +/-000	0.040

Table 2 Comparison of history among the group of oats and flaxseed group.

Background	Oats (n)	Flaxseed (n)	p*
Gender			
Male	9	12	0.213
Female	6	3	
History of hypercholesterolemia			0.341
Yes	3	5	
NO	12	10	
History of hypertriglyceridaemia			0.500
Yes	8	9	
NO	7	6	
History of Smoking			0.064
Yes	7	13	
NO	8	3	
Degree of obesity			0.362
Normal	3	1	
Overweight	4	3	
Obesity	8	9	
Severe obesity	0	2	

* Fisher's exact test

Confounding variables diet and exercise were monitored. In the case of the exercises there were no significant differences since in both cases it was low-impact exercise, unchanged during the study. For the degree of obesity both groups were similar, with no significant difference ($p = 0.362$) (Table 7), likewise the average Kcal amount consumed by the volunteer was similar (2786.20 ± 970.01 Kcal from group 1 (oats) against 2762.00 ± 1052.64 kcal of group 2 (flaxseed) with $p = 0.948$). However, it was found that in both groups there was a significantly reduction in weight (Tables 8 and 9).

Discussion

In the present study, the use of oats and flaxseeds to lower triglyceride levels were not effective, so you should accept the null hypothesis. It is likely that the sample size is influenced decisively, since triglyceride levels of the flaxseed group fell by almost 15% from their original levels, and if this trend continued -with an increase of volunteers. We would have obtained more encouraging results.

On the other hand, it's worth pointing out that in both groups there was significant reduction in weight, without a strict diet or exercise routine. Also, both treatments were well tolerated and

Table 3 Comparison of lipid levels at the beginning and end of treatment oatmeal.

Profile	Before Mean (SD) medium	After Mean (SD) medium	p*
Triglycerides mg/dL	298.80 +/-114.60269	307.20 +/-126.95304	0.798
Cholesterol mg/dL	205.20 +/-37.28209.00	204.27 +/-28.40209.00	0.798
LDL mg/dL	104.08 +/-24.82111.00	110.00 +/-26.83105.00	0.712
HDL mg/dL	44.80 +/-9.0943.00	43.80 +/-8.2742.00	0.674

* Test for paired samples ranges of Wilcoxon

Table 4 Comparison of lipid levels at the beginning and end with the treatment of flaxseed.

Profile	Before Mean (SD) medium	After Mean (SD) medium	p*
Triglycerides mg/dL	351.62 +/- 155.17282.00	303.23 +/-117.40289.00	0.152
Cholesterol mg/dL	215.85 +/-39.85204.00	200.62 +/- 32.87193.00	0.234
LDL mg/dL	101.23 +/-34.2393.00	106.62 +/-27.47103.00	0.944
HDL mg/dL	37.86 +/-6.4637.50	39.92 +/-5.5642.00	0.588

* Test for paired samples ranges of Wilcoxon

Table 5 Comparison of levels of renal and liver function at the beginning and end, with oat.

Profile	Before Mean (SD) medium	After Mean (SD) medium	p*
Direct bilirubin mg/dl	0.16 +/- 0.08 0.16	0.15 +/-0.070.14	0.281
ALT mU/ml	21.40 +/-6.820	21.40 +/-7.122	0.907
Urea mg/dL	28.46 +/-4.6627.80	28.66 +/-5.1130	0.925
Glucose mg/dl	123.53 +/-76.69100	143.47 +/-157.35108	0.865

*Test for paired samples ranges of Wilcoxon

Table 6 Comparison of levels of liver and kidney function at the beginning and end with flaxseed.

Profile	Before Mean (SD) medium	After Mean (SD) medium	p*
Direct bilirubin mg/dl	0.17 +/- 0.06 0.18	0.23 +/-0.20 0.19	0.460
ALT mU/ml	29.31 +/-10.04 26	25.08 +/-6.9 23	0.154
Urea mg/dL	28.98 +/-6.61 30	30.01 +/-8.78 27.90	0.657
Glucose mg/dl	117.69 +/-59.20 95	117.92 +/-57.83 100	0.674

* Test for paired samples ranges of Wilcoxon

Table 7 Distribution of patients according to their weight.

		Treatment		Total
		Treatment with oat	Treatment with flaxseed	
Degree of Obesity	Normal	3	1	4
	Over weight	4	3	7
	Obesity	8	9	17
	Severe obesity	0	2	2
Total		15	15	30

safe so that only 2 volunteer's flaxseed group had to suspend the treatment for less serious gastrointestinal complaints in which there were no complications following their suspension. They were safe as no any alteration of the liver or renal function so they can be administered without substantial risk found.

So far, the results of other studies on the use of oatmeal in reducing serum triglyceride levels have been poor; while studies in rats were promising, they were, however, frankly disappointing in humans. Nevertheless, natural fibers both oats and flaxseed should continue being recommended as part of the management strategy for patients with risk factors and to a healthy diet without forgetting its effect on weight and the bowel habits.

Despite the results and explanations, we believe that we must continue with more research on the use of oats and other natural

Table 8 Effect on weight with oats.

	Antes Media +/- DE Mediana	Después Media +/- DE Mediana	p*
Weight (kg)	85.29 +/-18.40 83.20	84.42 +/-17.86 83.7	0.007
BMI	31.42 +/-5.47 31.49	31.15 +/-5.32 31.56	0.006

* Test for paired samples combined ranges of Wilcoxon

Table 9 Effect on weight with flaxseed.

	Before Mean +/- SD Medium	After Mean +/- SD Medium	p*
Weight (kg)	89.76 +/-13.80 90.50	89.46 +/-13.60 90.00	0.043
BMI	32.26 +/-5.1 32.31	31.37 +/-5.74 31.73	0.020

* Test for paired samples combined ranges of Wilcoxon

fibers in different doses and times, as with a larger sample size because compared to drugs already established for management of dyslipidemia, the natural fibers are much cheaper and with poor side effects and also can be used even in the absence of cardiovascular risk factors.

Conclusions

The oral use of oats is safe and well tolerated but was ineffective for lowering serum triglyceride levels; similarly, the use of flaxseed in volunteers with hypertriglyceridaemia wasn't effective either. However, it was found that in both treatments there was a significantly weight loss.

Acknowledgements

We appreciate the participation of relevant form to:

- Maria Margarita Flores Castillo, Nutritionist, General

Hospital No. 47

- Francisco Hernández Balderas, Translation Assistant.

References

- 1 Munguía-Miranda C, Sánchez-Barrera RG, Hernández-Saavedra D, Cruz-López C (2008) Prevalencia de dislipidemias en una población de sujetos en apariencia sanos y su relación con la resistencia a la insulina. *Salud Publica Mex* 50: 375-382.
- 2 Martínez-Hernández AF, Chávez-Aguirre R (2007) Prevalencia y comorbilidad de dislipidemias en el primer nivel de atención. *Rev Med Inst Mex Seguro Soc* 45: 469-475.
- 3 Camacho-Hernández R, Corona-Muñiz I, Vázquez-Martínez JL, Martínez-Rodríguez F, Escobedo-de la Peña P (1995) Risk factors for ischemic heart disease in Mexico: a case control study. *Arch Inst Cardiol Mex* 65: 315-322.
- 4 Posadas-Romero C, Sepúlveda J, Tapia-Conyer R, Magos C, Cardoso-Magaña GC, et al. (1992) Valores de Colesterol Sérico en la Población Mexicana. *Salud Pública de México* 34: 157-167.
- 5 Verdejo J, Llamas G, Manejo de la (2000) Aterosclerosis en la Fase Crónica. *Arch Inst Cardiol Méx* 70: 83-90.
- 6 Sacks FM, Pfeffer MA, Moye LA, Rouleau JL, Rutherford JD, et al. (1996) The Effect of Pravastatin on Coronary Events after Myocardial Infarction in Patients with Average Cholesterol Levels. *N Engl J Med* 335: 1001-1009.
- 7 Ginsberg HN, Elam MB, Lovato LC, Crouse JR, Leiter LA, et al. (2010) Effects of combination lipid therapy in type 2 diabetes mellitus. *N Engl J Med* 362: 1563-1574.
- 8 Valdivielso P, Pintó X, Mateos-Gallegos R, Masana L, Alvarez-Salas L, et al. (2011) Características clínicas de los pacientes con hipertrigliceridemia remitidos a las Unidades de Lípidos: registro de hipertrigliceridemias de la Sociedad Española de Arteriosclerosis. *Med Clin (Barc)* 136: 231-238.
- 9 Brunzell J (2007) Todo sobre hipertrigliceridemia. *N Engl J Med* 357: 1009-1017.
- 10 Hernández-Pérez F, Herrera-Arellano A (2011) Tratamiento de la hipercolesterolemia con *Hibiscus sabdariffa* Ensayo clínico aleatorizado controlado. *Rev Med del Inst Mexi del Seg Social* 49: 469-480.
- 11 Ros E, Laguna J (2006) Tratamiento de la hipertrigliceridemia: fibratos frente a ácidos grasos omega-3. *Rev Esp Cardiol Supl* 6: 52D-61D.
- 12 Peñafiel D, Guatemal W (2010) Prevalencia de dislipidemias y sus factores de riesgo en adultos que acuden al Centro de Salud No. 1 de la ciudad de Ibarra, provincia de Imbabura octubre (Tesis de grado previo). Ibarra, Ecuador. Tesis de grado previo a la obtención del título de la licenciatura en nutrición y salud comunitaria.
- 13 Cortes García JC, Sosa Manriquez LF, y col (2010) Efecto de la avena sativa sobre los lípidos plasmáticos en habitantes de Nueva Italia y Lombardía sin restricción dietética. Morelia, Michoacán. Tesis para obtener el título de químico fármaco-biólogo de Patsy Nadhire Murillo Barcenas.
- 14 Morón MCh, Infante B, Ávila AV, García OE, Liuzzi JP (2010) Efecto del consumo de dietas con avena y carotas negras sobre el perfil lipídico en un modelo experimental en ratas. *Revista del Instituto Nacional de Higiene "Rafael Rangel"* 41: 31-37.
- 15 Glore SR, Van Treeck D, Knehans Aw, Guild M (1994) Soluble fiber and serum lipids: a literature review. *J Am Diet Assoc* 94: 425-436.
- 16 Kris-Etherton P, Krummel D, Russell ME, Dreon D, Mackey S, et al. (1988) The effect of diet and plasma lipids, lipoproteins and coronary heart disease. *J Am Diet assoc* 88: 1373-1400.
- 17 Anderson JW, Hanna TJ (1999) Impact of non-digestible carbohydrates on serum lipoprotein and risk for cardiovascular disease. *J Nutr* 129: 1457S -1466S.
- 18 Haskell W, Spiller G, Jensen C, Ellis B, Gates J (1992) Role of water-soluble dietary fiber in the management of elevated cholesterol in healthy subjects. *Am J Cardiol* 69: 433-439.
- 19 Brown L, Rosner B, Willett WW, Sacks FM (1999) Cholesterol lowering effects of dietary fiber: a meta-analysis. *Am J Clin Nutr* 69: 30-42.
- 20 Moya ME, Galussi AA, Reinoso DP, Soldá GI (2002) Caracterización varietal de avena sativa l. y avena byzantina c. koch por electroforesis de aveninas y proteínas totales en semillas. *Revista Científica Agropecuaria* 6: 41-47.
- 21 Othman RA, Moghadasian MH, Jones PJ (2004) Cholesterol-lowering effects of oat β -glucan. *Nutrition Reviews* 69: 299-309.
- 22 Kerckhoffs DA, Brouns F, Hornstra G, Mensink RP (2002) Effects on the Human Serum Lipoprotein Profile of beta-Glucan, Soy Protein and Isoflavones, Plant Sterols and Stanols, Garlic and Tocotrienols. *J Nutr* 132: 2494-2505.
- 23 Ruiz-Félix E, Mejía-Rodríguez O, Herrera-Abarca A, Cortés-García JC (2011) Consumo de avena (*Avena sativa*) y prevención primaria de la dislipidemia en adultos sin restricción dietética *Aten Fam* 18: 35-37.
- 24 Naumann E, Rees AB, Önnings G, Öste R, Wydra M, et al. (2006) Beta-Glucan incorporated into a fruit drink effectively lowers serum LDL-cholesterol concentrations. *Am J Clin Nutr* 83: 601-605.
- 25 Brown L, Rosner B, Willett WW, Sack FM (1999) Cholesterol-lowering effects of dietary fiber: a meta-analysis. *Am J Clin Nutr* 69: 30-42.
- 26 Kerckhoffs DA, Hornstra G, Mensink RP (2003) Cholesterol-lowering effect of beta-glucan from oat bran in mildly hypercholesterolemic subjects may decrease when beta-glucan is incorporated into bread and cookies. *Am J Clin Nutr* 78: 221-227.
- 27 Björklund M, Van Rees A, Mensink RP, Önnings G (2005) Changes in serum lipids and postprandial glucose and insulin concentrations after consumption of beverages with beta-glucans from oats or barley: a randomised dose-controlled trial. *Eur J Clin Nutr* 59: 1272-1281.
- 28 Karmally W, Montez MG, Palmas W, Branstetter A, Ramakrishnan R, et al. (2005) Cholesterol-lowering benefits of oat-containing cereal in Hispanic Americans. *J Am Diet Assoc* 105: 967-970.
- 29 Maki KC, Davidson MH, Ingram KA, Veith PE, Bell M, et al. (2003) Lipid responses to consumption of a beta-glucan containing ready-to-eat cereal in children and adolescents with mild-to-moderate primary hypercholesterolemia. *Nutrition Research* 23: 1527-1535.
- 30 Lundin KE, Nilsen EM, Scott HG, Løberg EM, Goen A, et al. (2003) Oats induced villous atrophy in coeliac disease. *Nutrition Research* 23: 1527-1535.